


Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

A 99.9
E 76224
Cop. 2

1979

Northeastern Forest Experiment Station



FOREST SERVICE, U.S. DEPARTMENT OF AGRICULTURE, 370 REED ROAD, BROOMALL, PA. 19008

PRELIMINARY TEST OF BOOBY-TRAPPING FOR CONTROL OF TWO CERAMBYCIDS

—JIMMY R. GALFORD
Research Entomologist
Northeastern Forest Experiment Station,
Delaware, Ohio

FOREST SERVICE
CURRENT SERIALS
NOV 5 1979
U.S. DEPARTMENT OF AGRICULTURE
NATIONAL LIBRARY OF MEDICINE

ODC 414.11:453-145.7 x 19.88

Abstract. Small aluminum "boats" containing cotton saturated with lindane were glued to elytra of female red oak borers and locust borers and the beetles released into cages or onto trees. Males attempting to mate with booby-trapped females contacted the insecticide and died. However, females also became contaminated in mating attempts and soon died.

Keywords: red oak borer, locust borer, insecticide

INTRODUCTION

The idea of using "booby-trapping" for insect control was proposed by Smith (1963), Whitten and Norris (1967), and Morgan (1967). Masner et al. (1968) proposed a similar technique. The idea of booby-trapping is to release females or males that carry insecticide or chemosterilant and thus kill or sterilize potential mates of the released insects.

For insecticide treatment, it is necessary to have an insecticide-resistant strain of insects so that the treatment does not kill those carrying the booby-traps. Developing resistant strains can usually be accomplished within a few generations; however, the task is time

consuming with insects that have long life cycles.

The red oak borer, *Enaphalodes rufulus* (Haldeman) and locust borer *Megacyllene robiniae* (Forster), both primary borers, have long life cycles. The red oak borer has a 2-year life cycle with an economically important generation only in odd years throughout most of the beetles' range (Hay 1974). The locust borer has a 1-year life cycle. While both of these species can be reared artificially, a generation requires 7 to 8 months for the red oak borer (Galford 1974) and about 3 months for the locust borer (Wollerman 1969).

However, both cerambycids have potential for control by booby-trapping, because peak

male emergence occurs 1 to 2 weeks before peak female emergence.

The red oak borer is a prime candidate for booby-trapping because adults occur only in odd-numbered years. Adult activity lasts only about 6 weeks and average populations are 12 to 24 beetles per acre. Also, the adults are large and easily handled for affixing booby-traps.

This paper reports on preliminary studies with female beetles carrying insecticide-treated booby-traps.

MATERIALS AND METHODS

The red oak borers used in this study were artificially reared (Galford 1974). The locust borers were collected in late September from goldenrod flowers, *Solidago* spp.,. Female beetles were cooled in a refrigerator to facilitate their handling. Small pieces of cotton wadding were cut to fit inside small aluminum combustion boats (#29-410, Coleman Instruments Corp. Maywood, IL).¹ The cotton was glued inside the boats with contact cement. Next, the bottoms of the boats were dipped in contact cement and they were laid on their sides to dry until the cement became tacky.

The beetles were removed from the refrigerator and the boats were glued onto the elytra just behind the thorax. The beetles were then released in well-ventilated containers until the glue dried.

An eye dropper was used to saturate the cotton with 0.1 ml of a 20 percent lindane solution. Ten female red oak borers and six female locust borers were prepared. The red oak borers were tested as follows: Each female red oak borer was released singly in 1-liter plastic bucket, the sides and bottom of which were lined with filter paper to enable the beetles to crawl around without falling.

¹The use of trade, firm, or corporation names in this publication is for the information and convenience of the reader. Such use does not constitute an official endorsement or approval by the U.S. Department of Agriculture or the Forest Service of any product or service to the exclusion of others that may be suitable.

Three male beetles were introduced into each bucket and a perforated lid was placed on top. After 24 hours beetle mortality was recorded.

Treated female locust borers were released singly on trees in a black locust plantation. Small trees were selected so that the treated females could be easily observed and any males contacting them could be collected. Any male beetles that made contact with or attempted to mate with treated females were collected and placed in individual jars. After 24 hours male mortality was recorded.

RESULTS AND DISCUSSION

All 30 of the male red oak borers were dead or near death after 24 hours; however, so were the females. The aluminum boats were intended to prevent the insecticide from contacting the females. However, male beetles attempting to mate contaminated the booby-trapped females with insecticide. Not all the locust borer males attempted to mate with the females they encountered. When their antennae contacted the boats before the body of the female, they were apparently repelled by the odor of the insecticide and moved away. The boats were apparently too large for the locust borer females and greatly reduced the body area a male could contact. This is important because males recognize females by touch.

Eleven male locust borers that made contact with the six released females were collected. Nine of the 11 males died within 24 hours. The other two males appeared unaffected. Three other males made contact with the females but escaped. All six treated females fell from the trees within 30 minutes and soon died.

While the test results show that females can be successfully booby-trapped, it also demonstrates the necessity of using insecticide-resistant strains. If effective chemosterilants for cerambycids could be found, they would greatly enhance control by booby-trapping.

LITERATURE CITED

- Galford, J. R.
1974. **Some physiological effects of temperature on artificially reared red oak borers.** J. Econ. Entomol. 67:709-710.
- Hay, C. J.
1974. **Survival and mortality of red oak borer larvae on black, scarlet, and northern red oak in eastern Kentucky.** Ann. Entomol. Soc. Am. 67(6):981-986.
- Masner, P., K. Slama and V. Landa
1968. **Sexually spread insect sterility induced by the analogues of juvenile hormone.** Nature 219:395-396.
- Morgan, P. B.
1967. **Booby-trapped female house flies or sterilant carriers.** J. Econ. Entomol. 60:612-613.
- Smith, C. N.
1963. **Prospects for vector control through sterilization procedures.** Bull. World Health Organ. 29(suppl.):99-106.
- Whitten, M. J. and K. R. Norris
1967. **"Booby-trapping" as an alternative to sterile males for insect control.** Nature 216: 1136.
- Wollerman, E. H., C. Adams and G. C. Heaton
1969. **Continuous laboratory culture of the locust borer, *Megacyllene robiniae*.** Ann. Entomol. Soc. Am. 62:647-649.



Use Pesticides Safely
FOLLOW THE LABEL

U.S. DEPARTMENT OF AGRICULTURE

Caution about Pesticides

This publication reports research involving pesticides. It does not contain recommendations for their use, nor does it imply that the uses discussed here have been registered. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife—if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended practices for the disposal of surplus pesticides and pesticide containers.
